

Are you ready for the next step?



You have a lot to consider when it comes to your experience with ovarian cancer. Whether you are undergoing treatment or the cancer has responded to treatment, understanding your next steps and how ZEJULA may fit in is important. That means having an open conversation with your doctor about all the options.

What I would like to share

➔ Use this first section to help you decide what you want to share with your doctor about your goals and what is important to you.

The challenges I have managing cancer right now are



My treatment goals include



My personal goals (the things that are important to me) are



What I would like to ask

➔ Now use this section to determine what questions you have about treatment options and if ZEJULA may be right for you.

☑ Check the questions you want to ask at your next appointment (circle the top 3).

- | | |
|--|---|
| <input type="checkbox"/> What else should I consider when we are deciding what to do next? | <input type="checkbox"/> How does ZEJULA work? |
| <input type="checkbox"/> When would ZEJULA be an option for me? | <input type="checkbox"/> How do I take ZEJULA? |
| <input type="checkbox"/> How might ZEJULA be part of my treatment plan? | <input type="checkbox"/> How will I know if ZEJULA is working? |
| <input type="checkbox"/> What is maintenance treatment? | <input type="checkbox"/> How will I know if my dose needs to be adjusted? |
| <input type="checkbox"/> Does my <i>BRCA</i> status matter when we are deciding if ZEJULA is an option for me? | <input type="checkbox"/> What are important side effects I should know about? |
| | <input type="checkbox"/> If I have side effects, what can I do about them? |

Indication(s)

ZEJULA is a prescription medicine used for the:

- maintenance treatment of adults with advanced ovarian cancer, fallopian tube cancer, or primary peritoneal cancer. ZEJULA is used after the cancer has responded (complete or partial response) to treatment with platinum-based chemotherapy
- maintenance treatment of adults with a certain type of inherited (germline) abnormal *BRCA* gene with ovarian cancer, fallopian tube cancer, or primary peritoneal cancer that comes back. ZEJULA is used after the cancer has responded (complete or partial response) to treatment with platinum-based chemotherapy. Your healthcare provider will perform a test to make sure that ZEJULA is right for you.

It is not known if ZEJULA is safe and effective in children.

Please see Important Safety Information on back and accompanying Prescribing Information, also available at ZEJULA.com.

Important Safety Information



ZEJULA may cause serious side effects, including:

Bone marrow problems called Myelodysplastic Syndrome (MDS) or a type of blood cancer called Acute Myeloid Leukemia (AML). Some people who have ovarian cancer and who have received previous treatment with chemotherapy or certain other medicines for their cancer have developed MDS or AML during treatment with ZEJULA. MDS or AML may lead to death.

Symptoms of low blood cell counts (low red blood cells, low white blood cells, and low platelets) are common during treatment with ZEJULA. They can be a sign of serious bone marrow problems, including MDS or AML. These symptoms may include the following:

- Weakness
- Feeling tired
- Weight loss
- Frequent infections
- Fever
- Shortness of breath
- Blood in urine or stool
- Bruising or bleeding more easily

Your doctor will do blood tests to check your blood cell counts before treatment with ZEJULA. You will be tested weekly for the first month of treatment with ZEJULA, monthly for the next 11 months of treatment, and as needed afterward.

High blood pressure is common during treatment with ZEJULA, and it can become serious. Your doctor will check your blood pressure and heart rate at least weekly for the first two months, then monthly for the first year, and as needed thereafter during your treatment with ZEJULA.

Posterior reversible encephalopathy syndrome (PRES) is a condition that affects the brain and may happen during treatment with ZEJULA. If you have headache, vision changes, confusion, or seizure, with or without high blood pressure, please contact your doctor.

Before starting to take ZEJULA, tell your doctor about all of your medical conditions, including if you:

- Have heart problems
- Have liver problems
- Have high blood pressure
- Are allergic to FD&C Yellow No. 5 (tartrazine) or aspirin. ZEJULA capsules contain tartrazine, which may cause allergic-type reactions (including bronchial asthma) in certain people, especially people who also have an allergy to aspirin
- Are pregnant or plan to become pregnant. ZEJULA may harm an unborn baby and may cause loss of pregnancy (miscarriage)
 - If you are able to become pregnant, you should use effective birth control (contraception) during treatment with ZEJULA and for 6 months after taking the last dose of ZEJULA
 - If you are able to become pregnant, your doctor may perform a pregnancy test before you start treatment with ZEJULA
 - You should tell your doctor right away if you become pregnant
- Are breastfeeding or plan to breastfeed
 - ZEJULA may harm your baby. You should not breastfeed your baby during treatment with ZEJULA and for 1 month after taking the last dose of ZEJULA

Tell your doctor about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

The most common side effects of ZEJULA include the following:

- Heart not beating regularly
- Nausea
- Constipation
- Vomiting
- Pain in the stomach area
- Mouth sores
- Diarrhea
- Indigestion or heartburn
- Dry mouth
- Tiredness
- Loss of appetite
- Urinary tract infection
- Changes in liver function or other blood tests
- Pain in your muscles and back
- Headache
- Dizziness
- Change in the way food tastes
- Trouble sleeping
- Anxiety
- Sore throat
- Shortness of breath
- Cough
- Rash
- Changes in the amount or color of your urine

If you have certain side effects, then your doctor may change your dose of ZEJULA, temporarily stop, or permanently stop treatment with ZEJULA.

These are not all the possible side effects of ZEJULA. For more information, ask your doctor or pharmacist. Call your doctor for medical advice about side effects. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

Please see full Prescribing Information.

Trademarks are owned by or licensed to the GSK group of companies.



©2023 GSK or licensor.
NRPWCNT230006 March 2023
Produced in USA.